



## Short communication

# Dose-response analysis indicating time-dependent neurotoxicity caused by organic and inorganic mercury—Implications for toxic effects in the developing brain



Julia Pletz<sup>a,\*</sup>, Francisco Sánchez-Bayo<sup>b</sup>, Henk A. Tennekens<sup>a</sup>

<sup>a</sup> Experimental Toxicology Services (ETS) Nederland BV, Frankensteeg 4, 7201 KN Zutphen, The Netherlands

<sup>b</sup> Dept. Plant and Food Sciences, Faculty of Agriculture & Environment, The University of Sydney, 1 Central Avenue, C81-ATP, Eveleigh, NSW 2015, Australia

## ARTICLE INFO

## Article history:

Received 27 January 2016

Received in revised form 19 February 2016

Accepted 25 February 2016

Available online 2 March 2016

## Keywords:

Dose-response modelling

Toxicodynamics

Mercury

Neurotoxicity

Risk assessment

## ABSTRACT

A latency period preceding neurotoxicity is a common characteristic in the dose-response relationship induced by organic mercury. Latency periods have typically been observed with genotoxicants in carcinogenesis, with cancer being manifested a long time after the initiating event. These observations indicate that even a very small dose may cause extensive adverse effects later in life, so the toxicity of the genotoxic compound is dose and time-dependent. In children, methylmercury exposure during pregnancy (in utero) has been associated with delays in reaching developmental milestones (e.g., age at first walking) and decreases in intelligence, increasing in severity with increasing exposure. Ethylmercury exposure from thimerosal in some vaccines has been associated, in some studies, with autism and other neurological disorders in children. In this paper, we have examined whether dose-response data from *in vitro* and *in vivo* organic mercury toxicity studies fit the Druckrey-Küpfmüller equation  $c \cdot t^n = \text{constant}$  ( $c$  = exposure concentration,  $t$  = latency period), first established for genotoxic carcinogens, and whether or not irreversible effects are enhanced by time of exposure ( $n \geq 1$ ), or else toxic effects are dose-dependent while time has only minor influence on the adverse outcome ( $n < 1$ ). The mode of action underlying time-dependent toxicity is irreversible binding to critical receptors causing adverse and cumulative effects. The results indicate that the Druckrey-Küpfmüller equation describes well the dose-response characteristics of organic mercury induced neurotoxic effects. This amounts to a paradigm shift in chemical risk assessment of mercurial compounds and highlights that it is vital to perform toxicity testing geared to investigate time-dependent effects.

© 2016 Elsevier Ireland Ltd. All rights reserved.

## 1. Introduction

Organic mercury induced neurotoxicity has typically been observed after a preceding latency period. Even severe and fatal ethylmercury intoxications in humans featured a latency period between cessation of exposure and onset of first symptoms of 10 days to 7 weeks (Cinca et al., 1980; Magos, 2001). For methylmercury, latencies in intoxications in Iraq and Japan ranged from weeks to more than a year (Bakir et al., 1973; National Research Council, 2000; Weiss et al., 2002) and effects were proceeding even after exposure had ended 20–30 years before (Rice and Barone, 2000). When monkeys were exposed to low levels of methylmercury during their developmental phase,

neurotoxicity appeared after several years (Rice, 1996). Latency periods have typically been observed with genotoxicants in carcinogenesis, with cancer being manifested a long time after the initiating event. These observations indicate that even a very small dose may cause extensive adverse effects later in life, so the toxicity of the genotoxic compound is dose and time-dependent.

Methylmercury is widely distributed throughout the environment, particularly in estuarine and marine sediments (Bryan and Langston, 1992; Compeau and Bartha, 1985; Morel et al., 1998) and accumulates in fish and birds (Greichus et al., 1973; Harris et al., 2007; Henny et al., 2005; Houserová et al., 2007; Lam et al., 2005; Polak-Juszczak, 2012; Wren, 1986). Therefore, people are likely to be continuously exposed to small amounts of methylmercury through consumption of contaminated food (Chan et al., 2010; Lin et al., 2012). Ethylmercury is used as preservative in vaccines that may be administered to pregnant women. Toxicokinetic evidence confirms that alkyl mercury compounds cross the placental barrier (Aschner and Clarkson, 1988; Bridges and Zalups, 2005; Dórea,

\* Corresponding author. Present address: Feldbergstrasse 92, 4057 Basel, Switzerland.

E-mail addresses: [julia.pletz@yahoo.co.uk](mailto:julia.pletz@yahoo.co.uk) (J. Pletz), [sanchezbayo@mac.com](mailto:sanchezbayo@mac.com) (F. Sánchez-Bayo), [info@toxicology.nl](mailto:info@toxicology.nl) (H.A. Tennekens).

2011; Dórea et al., 2013; Geier et al., 2007; Kajiwara et al., 1996; Kerper et al., 1992) and the blood-brain barrier (Apostoli et al., 2006; National Research Council, 2000).

A myriad of scientific papers have been published on the toxicology of inorganic and organic mercury compounds. However, limited data are available for quantitative analysis of time-dependent toxicity. Most animal studies did not examine effects observed at different time points over extended exposure periods. Studies of human poisoning cases widely observed a latency period but an accurate estimation of dose or concentration at the target site is difficult to obtain. Data on Iraqi poisonings presented by Bakir et al. (1973) outlined the onset of symptoms after a mean latency period in relation to mercury concentrations in the blood of intoxicated individuals. These blood samples were obtained up to 115 days after the onset of symptoms. However, animal studies demonstrate a great divergence between blood and brain concentrations after exposure has ceased, suggesting that mercury persists in the brain while clearance from the blood occurs faster (Burbacher et al., 2005; Evans et al., 1977; Magos, 2001; Vahter et al., 1995). Therefore, blood concentrations taken a long time after exposure has ceased may not be a good parameter for assessing dose- and time-dependent neurotoxicity.

## 2. Dose and time-dependent toxicity of alkyl mercury compounds: data analysis

Wobeser et al. (1976) published data on the relationship between methylmercury dose and the occurrence of neurotoxic clinical signs in mink. A latency period was observed in all animals exhibiting toxicity. The higher the administered dose the earlier adverse effects appeared. When examining the total dose administered to the animals in the different dose groups it becomes apparent that low doses given continuously over a long

period until the occurrence of ataxia add up to a lower total dose than the total dose from higher doses over a shorter period of time (Table 1). Decreasing total doses and increasing adverse effect induction times could reflect dose and time dependencies according to the Druckrey-Küpfmüller equation (Druckrey and Küpfmüller, 1949):

$$dt^n = \text{constant} \quad (1)$$

where the exponent  $n > 1$  can be regarded as an exposure-time-reinforcement factor (Tennekes and Sánchez-Bayo, 2013). A dose-time-effect analysis of the ataxia data provided by Wobeser et al. (1976) results in a value of  $n = 1.5$  ( $r^2 = 0.93$ ), as illustrated in Fig. 1.


Another dataset confirming that neurotoxic effects from methylmercury exposure are reinforced over time was established using data obtained by Grant (1973) in cats (Table 2). In this case, the Druckrey-Küpfmüller equation described the dose-response relationship with  $n = 2.9$  ( $r^2 = 0.71$ ). This high  $n$ -value may be related to the observation that cats were particularly sensitive to methylmercury exposure (Charbonneau et al., 1976). A latency period of at least 59 days was noted and clinical signs observed included ataxia and convulsions. Interestingly, when brain concentrations are correlated to the latency period in the onset of clinical signs a lower  $r^2$  value is obtained. Mercury is known to accumulate in specific brain regions which may not have been taken into account when brain concentrations were determined.

For ethylmercury-induced effects, the Druckrey-Küpfmüller equation describes the dose-response relationship in human cortical neurotoxicity observed by Baskin et al. (2003), (Table 3) with an exposure-time-reinforcement factor of 2.81 ( $r^2 = 0.8$ ), (Fig. 2) for concentrations ranging from 1 to 250  $\mu\text{M}$  at time points between 2 and 24 h. The extremely high value of  $n$  obtained in this case indicates pronounced reinforcement of toxicity by exposure time.

**Table 1**

Effect of dose on onset of methylmercury-induced ataxia in mink (after data from Wobeser et al., 1976).

Dose (d) in feed (ppm)	Median time (t50) to onset of ataxia (days)	Total dose (ppm) = d · t50
1.8	69	124.2
4.8	30.5	146.4
8.3	20	166
15	17.75	266.25



**Fig. 1.** Onset of ataxia in mink exposed to various doses of methylmercury in feed over time.

**Table 2**

Effect of methylmercury dose on the onset of neurotoxic signs in cats (after data from Grant,1973).

Daily dose (d), (mg/kg)	Latency period of clinical signs (t), (days)	Total dose (mg/kg)= d·t
0.34	71	24.14
0.44	63	27.72
0.47	68	31.96
0.55	59	32.45

**Table 3**

Effect of exposure concentration on latency periods of neurotoxicity in human cortical neurons exposed to ethylmercury (after data from Baskin et al., 2003).


Concentration ( $\mu\text{M}$ )	Time (t) to neurotoxic effect (hours)	Total dose ( $\mu\text{M}$ ) = c·t
1	24	24
2	6	12
10	4	40
50	4	200
250	2	500

### 3. Pharmacokinetic and pharmacodynamic basis of mercury-induced time-reinforced neurotoxicity

Druckrey and Küpfmüller (1949) explained time-reinforced toxicity ( $n > 1$ ) of chemicals as a result of irreversible receptor binding associated with an irreversible effect. The Druckrey-Küpfmüller theorem was recently validated for a number of genotoxic and non-genotoxic compounds, including inorganic (e.g. selenium) and organic chemicals with specific mode of action such as pesticides (Tennekes and Sánchez-Bayo, 2013).

In the case of mercury, the actual toxicant in the central nervous systems (CNS) is thought to be the divalent mercuric ion ( $\text{Hg}^{2+}$ ) which is formed when organic mercury compounds such as methyl- and ethylmercury dealkylate. Once organic mercury compounds reach the brain tissue and dealkylate,  $\text{Hg}^{2+}$  gets trapped in the neurons, as it cannot permeate the blood-brain barrier. In mink and monkeys, inorganic mercury remains in the brain even months after organic mercury exposure has ended (Burbacher et al., 2005; Evans et al., 1977; Vahter et al., 1994, 1995; Wobeser et al., 1976). A recent review outlined evidence from human case studies, animal studies and modelling assessments estimating an inorganic mercury half-life in human brains of at

least five to 27 years (Rooney, 2014). The “one-way” only kinetic pathway and the continuous dietary exposure lead to  $\text{Hg}^{2+}$  accumulation in the brain tissues. Neurons and glia in the cerebellum and hippocampus appear to be primary targets of this ion (Lohren et al., 2015; Pedersen et al., 1999).  $\text{Hg}^{2+}$  has electron-sharing facilities that can result in formation of covalent attachment to sulfhydryl groups of proteins, and binding of mercury species to thiol groups in amino acids, intracellular enzymes and structural proteins. For example, cysteine, glutathione and metallothionein are common targets of inorganic mercury and methylmercury (Conner and Fowler, 1993; Krupp et al., 2008). Moreover, in an interaction with cysteine, methylmercury does not only form  $\text{Hg-S}$  covalent bonds but may also bind to nitrogen and to a smaller proportion to oxygen atoms (Krupp et al., 2008). Direct binding of mercuric ion to enzymes' active sites, if the sites contain sulfhydryl groups, may lead to inactivation of the enzymes, and the interaction with thiol-containing antioxidants such as glutathione can cause increased production of reactive oxygen species that damage lipids, protein and DNA (Ercal et al., 2001). It can be envisaged that mercury neurotoxicity would result from an autocatalytic process initiated by binding of mercuric ion to sulfhydryl groups of organic macromolecules.

**Fig. 2.** Latency period of toxicity in relation to exposure concentration in ethylmercury-induced neurotoxicity in human cortical neurons in vitro.

Autocatalytic processes have been observed in the Alzheimer's- and prion-disease-associated accumulation of amyloid fibrils, which has been linked to the induction of calcium influx and reactive oxygen species (ROS) leading to neuronal degeneration and oxidative stress (Ho et al., 2001; Kupfer et al., 2009; Zetterberg and Blennow, 2013). The accumulation of cytosolic calcium has been suggested as an early and pivotal event in causing neurotoxicity as it initiates all the other neurodegenerative events including generation of ROS and apoptosis (Ekinci et al., 1999, 2000). An increased  $\text{Ca}^{2+}$  influx has also been described as fundamental to mercury toxicity. Upon binding to glutathione and metallothionein, mercury initiates alterations in membrane permeability to calcium ions leading to oxidative stress, glutathione depletion and subsequent cell death (Conner and Fowler, 1993; Klaassen, 2008; Sarafian and Verity, 1991). Ethyl-, methyl- and inorganic mercury have all been demonstrated to be cytotoxic in differentiated human and rat neuron cultures at decreasing concentrations over an increasing exposure period (Fujimura and Usuki, 2012; Lohren et al., 2015; and as discussed Baskin et al., 2003).

The effect of interest here is developmental neurotoxicity. Methylmercury exposure to the developing brain compromises neuronal proliferation, migration and as a result neuronal differentiation, synaptogenesis, tightly regulated apoptosis, and other processes vital to the formation and functioning of the nervous system (for a detailed review see Rice and Barone, 2000; Rodier, 1995). The time window which encompasses the vulnerability of the brain to disturbances of all these processes is ample. In fact, it spans from the embryonic period until well into adulthood when a recapitulation of synaptogenesis is ongoing (Rice and Barone, 2000). The structures forming at the time of exposure play a decisive role on the extent of effects that show up later in life (Rodier, 1995). Therefore, a woman who is exposed during the first trimester of pregnancy may contract deficits or defects very different from those developed by someone who is exposed during the third trimester of pregnancy. However, with an extensive window of vulnerability (with respect to time and the amount of potential insults) and regular exposure (e.g. each portion of fish consumed by pregnant women at a frequency of one or two times a week), the potential of an insult of any of the critical processes such as neurogenesis, migration, differentiation or synaptogenesis is high. One developmental perturbation can initiate a cascade or autocatalytic processes of structural or functional changes which may be manifested as developmental delays or persistent deficits (Rice and Barone, 2000). Adverse effects may also appear staggered, with initial transient deficits and reversal when compensatory capability reduces during aging. Therefore, some effects may appear late in life when plasticity is declining or neurons are gradually lost. For that reason, compensation or neuronal plasticity only covers the neurotoxic effects for a certain period of time, while due to the limited capacity of the CNS to repair damage, such effects are usually irreversible and will manifest at some point in life. Although latently proceeding tissue injuries are common phenomena of toxicant-induced etiology, CNS plasticity is an organ-specific characteristic which naturally contributes to latencies in neurodegeneration.

#### 4. Conclusion

Experimental data in mammals (Tables 1 and 2) indicate that the Druckrey-Küpfmüller equation,  $dt^n = \text{constant}$ , first established for genotoxic carcinogens, also describes the dose-response characteristics of neurotoxic effects of organic mercury compounds. Exposure time reinforces the toxicant effect, since the total dose required to produce neurotoxicity decreases with increasing exposure time. Data outlined here and elsewhere (Rondeau et al., 2014; Tennekkes, 2010; Tennekkes and Sánchez-Bayo, 2013) demonstrate that time may reinforce toxic effects of compounds other

than genotoxic carcinogens. This amounts to a paradigm shift in chemical risk assessment. The decisive elements of such dose-response relationships are irreversible receptor binding associated with an irreversible effect, and latency periods for toxicity. It is vital to perform toxicity testing geared to investigate time-dependent effects. Also, it is crucial to further explore the cascading toxic effects in the brain to provide a greater insight into the biochemical and pathological processes ongoing while no clinical signs of neurodegeneration have yet become apparent. The analysis outlined here indicates that there may not be a safe level of mercury exposure. We are particularly concerned about exposure during pregnancy when considering the risk of adverse neurodevelopmental effects in the unborn child. Hence, the implications of these findings are the need to limit environmental concentrations of mercury and phase out the use of ethylmercury in vaccines.

#### Conflict of interest

None.

#### Acknowledgements

The authors wish to thank Prof. Janna Koppe, Em. Prof. of Neonatology of the University of Amsterdam, for her comments on the manuscript.

#### References

- Apostoli, P., Cornelis, R., Duffus, J., Hoet, P., Lison, D., Templeton, D., 2006. Environmental Health Criteria 234: Elemental Speciation in Human Health Risk Assessment. World Health Organisation.
- Aschner, M., Clarkson, T.W., 1988. Distribution of mercury 203 in pregnant rats and their fetuses following systemic infusions with thiol-containing amino acids and glutathione during late gestation. *Teratology* 38 (2), 145–155.
- Bakir, F., Damluji, S.F., Amin-Zaki, L., Murtadha, M., Khalidi, A., Al-Rawi, N.Y., Doherty, R.A., 1973. Methylmercury poisoning in Iraq. *Science* 230–240.
- Baskin, D.S., Ngo, H., Didenko, V.V., 2003. Thimerosal induces DNA breaks, caspase-3 activation, membrane damage, and cell death in cultured human neurons and fibroblasts. *Toxicol. Sci.* 74 (2), 361–368. doi:<http://dx.doi.org/10.1093/toxsci/kfg126>.
- Bridges, C.C., Zalups, R.K., 2005. Molecular and ionic mimicry of toxic metals. *Annu. Rev. Pharmacol. Toxicol.* 204 (3), 274–308. doi:<http://dx.doi.org/10.1146/annurev.pharmtox.33.1.545>.
- Bryan, G.W., Langston, W.J., 1992. Bioavailability: accumulation and effects of heavy metals in sediments with special reference to United Kingdom estuaries: a review. *Environ. Pollut.* 76, 89–131.
- Burbacher, T.M., Shen, D.D., Liberato, N., Grant, K.S., Cernichiari, E., Clarkson, T., 2005. Comparison of blood and brain mercury levels in infant monkeys exposed to methylmercury or vaccines containing thimerosal. *Environ. Health Perspect.* 113 (8), 1015–1021. doi:<http://dx.doi.org/10.1289/ehp.7712>.
- Chan, H.M., Scheuhammer, A.M., Ferran, A., Loupelle, C., Holloway, J., Weech, S., 2010. Impacts of mercury on freshwater fish-eating wildlife and humans. *Hum. Ecol. Risk Assess.* 9, 867–883.
- Charbonneau, S., Munro, I., Nera, E., Al, E., 1976. Chronic toxicity of methylmercury in the adult cat. Interim report. *Toxicology* 5, 337–349.
- Cinca, I., Dumitrescu, I., Onaca, P., Serbănescu, A., Nestorescu, B., 1980. Accidental ethyl mercury poisoning with nervous system, skeletal muscle, and myocardium injury. *J. Neurol. Neurosurg. Psychiatry* 43 (2), 143–149. doi:<http://dx.doi.org/10.1136/jnnp.43.2.143>.
- Compeau, G.C., Bartha, R., 1985. Sulfate-reducing bacteria: principal methylators of mercury in anoxic estuarine sediment. *Appl. Environ. Microbiol.* 50, 498–502.
- Conner, E.A., Fowler, B.A., 1993. Mechanisms of metal-induced nephropathy. In: Hook, J.B., Goldstein, R.S. (Eds.), *Toxicology of the Kidney*, second (edition) Raven Press, New York, pp. 437–457.
- Dórea, J.G., Farina, M., Rocha, J.B.T., 2013. Toxicity of ethylmercury (and thimerosal): a comparison with methylmercury. *J. Appl. Toxicol.* 33 (8), 700–711. doi:<http://dx.doi.org/10.1002/jat.2855>.
- Dórea, J.G., 2011. Integrating experimental (in vitro and in vivo) neurotoxicity studies of low-dose thimerosal relevant to vaccines. *Neurochem. Res.* 36 (6), 927–938. doi:<http://dx.doi.org/10.1007/s11064-011-0427-0>.
- Druckrey, H., Küpfmüller, K., (1949). *Dosis und Wirkung – Beiträge zur theoretischen Pharmakologie* (First). Freiburg im Breisgau, Germany: Cantor GmbH.
- Ekinci, F.J., Malik, K.U., Shea, T.B., 1999. Activation of the L voltage-sensitive calcium channel by mitogen-activated protein (MAP) kinase following exposure of neuronal cells to beta-amyloid. *J. Biol. Chem.* 274 (42), 30322–30327. doi:<http://dx.doi.org/10.1074/jbc.274.42.30322>.

- Ekinici, F.J., Linsley, M.D., Shea, T.B., 2000. Beta-amyloid-induced calcium influx induces apoptosis in culture by oxidative stress rather than tau phosphorylation. *Mol. Brain Res.* 76 (2), 389–395. doi:10.1016/S0169-328X(00)00025-5.
- Ercal, N., Gurer-Orhan, H., Aykin-Burns, N., 2001. Toxic metals and oxidative stress part I: mechanisms involved in metal induced oxidative damage. *Curr. Top. Med. Chem.* 1, 529–539.
- Evans, H.L., Garman, R.H., Weiss, B., 1977. Methylmercury: exposure duration and regional distribution as determinants of neurotoxicity in nonhuman primates. *Toxicol. Appl. Pharmacol.* 41, 15–33. doi:10.1016/0041-008X(77)90051-5.
- Fujimura, M., Usuki, F., 2012. Differing effects of toxicants (methylmercury, inorganic mercury, lead, amyloid beta, and rotenone) on cultured rat cerebrocortical neurons: differential expression of rho proteins associated with neurotoxicity. *Toxicol. Sci.* 126 (2), 506–514. doi:10.1093/toxsci/kfr352.
- Geier, D.A., Sykes, L.K., Geier, M.R., 2007. A review of thimerosal (merthiolate) and its ethylmercury breakdown product: specific historical considerations regarding safety and effectiveness. *J. Toxicol. Environ. Health B Crit. Rev.* 10 (8), 575–596. doi:10.1080/10937400701389875.
- Grant, C.A., 1973. Pathology of experimental methyl mercury intoxication: some problems of exposure and response. In: Miller, M.W., Clarkson, T.W. (Eds.), *Mercury, Mercurials and Mercaptans*. Charles C Thomas Publisher, Springfield, IL, pp. 294–312.
- Greichus, Y., Greichus, A., Emerick, R., 1973. Insecticides, polychlorinated biphenyls and mercury in wild cormorants, pelicans, their eggs: food and environment. *Bull. Environ. Contam. Toxicol.* 9, 321–328.
- Harris, R.C., Rudd, J.W., Amyot, M., Babiarz, C.L., Beaty, K.G., Blanchfield, P.J., et al., 2007. Whole-ecosystem study shows rapid fish-mercury response to changes in mercury deposition. *Proc. Natl. Acad. Sci. U. S. A.* 104, 16536–16591.
- Henny, C.J., Kaiser, J.L., Packard, H.A., Grove, R.A., Taf, M.R., 2005. Assessing mercury exposure and effects to American dippers in headwater streams near mining sites. *Ecotoxicology* 14, 709–725.
- Ho, P.I., Collins, S.C., Dhitavat, S., Ortiz, D., Ashline, D., Rogers, E., Shea, T.B., 2001. Homocysteine potentiates beta-amyloid neurotoxicity: role of oxidative stress. *J. Neurochem.* 78 (2), 249–253. doi:10.1046/j.1471-4159.2001.00384.x.
- Houserová, P., Kubán, V., Kráčmar, S., Sitko, J., 2007. Total mercury and mercury species in birds and fish in an aquatic ecosystem in the Czech Republic. *Environ. Pollut.* 145, 185–194.
- Kajiwara, Y., Yasutake, A., Adachi, T., Hirayama, K., 1996. Methylmercury transport across the placenta via neutral amino acid carrier. *Arch. Toxicol.* 70 (5), 310–314.
- Kerper, L.E., Ballatori, N., Clarkson, T.W., 1992. Methylmercury transport across the blood-brain barrier by amino acid carrier. *Am. J. Physiol.* 262 (5 Part 2), R761–765.
- Klaassen, C.D., 2008. Casarett and Doull's toxicology: the basic, seventh (edition) In: Klaassen, C.D. (Ed.), *Toxicology*, Vol. 12. McGraw-Hill, New York doi:10.1036/0071470514.
- Krupp, E.M., Milne, B.F., Mestrot, A., Meharg, A.A., Feldmann, J., 2008. Investigation into mercury bound to biothiols: structural identification using ESI-ion-trap MS and introduction of a method for their HPLC separation with simultaneous detection by ICP-MS and ESI-MS. *Anal. Bioanal. Chem.* 390 (7), 1753–1764. doi:10.1007/s00216-008-1927-x.
- Kupfer, L., Hinrichs, W., Groschup, M.H., 2009. Prion protein misfolding. *Curr. Mol. Med.* 9 (7), 826–835. doi:10.2174/156652409789105543.
- Lam, J.C.W., Tanabe, S., Lam, M.H.W., Lam, P.K.S., 2005. Risk to breeding success of waterbirds by contaminants in Hong Kong: evidence from trace elements in eggs. *Environ. Pollut.* 135, 481–490.
- Lin, Y., Vogt, R., Larssen, T., 2012. Environmental mercury in China: a review. *Environ. Toxicol. Chem.* 31, 2431–2444.
- Lohren, H., Blagojevic, L., Fitkau, R., Ebert, F., Schildknecht, S., Leist, M., Schwerdtle, T., 2015. Toxicity of organic and inorganic mercury species in differentiated human neurons and human astrocytes. *J. Trace Elem. Med. Biol.* 32, 200–208. doi:10.1016/j.jtemb.2015.06.008.
- Magos, L., 2001. Review on the toxicity of ethylmercury including its presence as a preservative in biological and pharmaceutical products. *J. Appl. Toxicol.* 21 (1), 1–5. doi:10.1002/jat.721.
- Morel, F.M.M., Kraepiel, A.M.L., Amyot, M., 1998. The chemical cycle and bioaccumulation of mercury. *Annu. Rev. Ecol. Syst.* 29, 543–566.
- National Research Council, 2000. *Toxicological Effects of Methylmercury*. National Academies Press, Washington, DC. <http://www.nap.edu/catalog/9899.html>.
- Pedersen, M.B., Hansen, J.C., Mulvad, G., Pedersen, H.S., Gregersen, M., Danscher, G., 1999. Mercury accumulations in brains from populations exposed to high and low dietary levels of methyl mercury. *Int. J. Circumpolar Health* 96–107.
- Polak-Juszczak, L., 2012. Bioaccumulation of mercury in the trophic chain of flatfish from the Baltic Sea. *Chemosphere* 89, 585–591.
- Rice, D.C., Barone, S., 2000. Critical periods of vulnerability for the developing nervous system: evidence from humans and animal models. *Environ. Health Perspect.* 108 (Suppl. 3), 511–533. doi:10.1289/ehp.00108s3511.
- Rice, D.C., 1996. Evidence for delayed neurotoxicity produced by methylmercury. *Neurotoxicology* 17, 583–596.
- Rodier, P.M., 1995. Developing brain as a target of toxicity. *Environ. Health Perspect.* 103 (Suppl. 6), 73–76. doi:10.1289/ehp.95103s673.
- Rondeau, G., Sánchez-Bayo, F., Tennekes, H.A., Decourtye, A., Ramírez-Romero, R., Desneux, N., 2014. Delayed and time-cumulative toxicity of imidacloprid in bees, ants and termites. *Sci. Rep.* 4, 1–8. doi:10.1038/srep05566.
- Rooney, J.P.K., 2014. The retention time of inorganic mercury in the brain—a systematic review of the evidence. *Toxicol. Appl. Pharmacol.* 274 (3), 425–435. doi:10.1016/j.taap.2013.12.011.
- Sarafian, T., Verity, M.A., 1991. Oxidative mechanisms underlying methyl mercury neurotoxicity. *Int. J. Dev. Neurosci.* 9 (2), 147–153.
- Tennekes, H.A., Sánchez-Bayo, F., 2013. The molecular basis of simple relationships between exposure concentration and toxic effects with time. *Toxicology* 309, 39–51. doi:10.1016/j.tox.2013.04.007.
- Tennekes, H.A., 2010. The significance of the Druckrey-Küpfmüller equation for risk assessment—the toxicity of neonicotinoid insecticides to arthropods is reinforced by exposure time. *Toxicology* 276 (1), 1–4. doi:10.1016/j.tox.2010.07.005.
- Vahter, M., Mottet, N.K., Friberg, L., Lind, B., Shen, D.D., Burbacher, T.M., 1994. Speciation of mercury in the primate blood and brain following long-term exposure to methyl mercury. *Toxicol. Appl. Pharmacol.* 124 (2), 221–229.
- Vahter, M.E., Mottet, N.K., Friberg, L.T., Lind, S.B., Charleston, J.S., Burbacher, T.M., 1995. Demethylation of methyl mercury in different brain sites of Macaca fascicularis monkeys during long-term subclinical methyl mercury exposure. *Toxicol. Appl. Pharmacol.* 134 (2), 273–284. doi:10.1006/taap.1995.1193.
- Weiss, B., Clarkson, T.W., Simon, W., 2002. Silent latency periods in methylmercury poisoning and in neurodegenerative disease. *Environ. Health Perspect.* 110 (Suppl. 5), 851–854. doi:10.1289/ehp.02110s5851.
- Wobeser, G., Nielsen, N.O., Schiefer, B., 1976. Mercury and mink II: experimental methyl mercury intoxication. *Can. J. Comp. Med.* 40 (January), 34–45.
- Wren, C.D., 1986. A review of metal accumulation and toxicity in wild mammals. *Environ. Res.* 40, 210–244.
- Zetterberg, H., Blennow, K., 2013. Biomarker evidence for uncoupling of amyloid build-up and toxicity in Alzheimer's disease. *Alzheimer's Dement.* 9 (4), 459–462. doi:10.1016/j.jalz.2012.07.002.